

45002-17



12/11/2014

1/37

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

Carolyn Miter  
Registration Specialist  
Albaugh, LLC  
PO Box 2127  
Valdosta, GA 31604

DEC 11 2014

Subject: Label Notification per PRN 98-10 – Add Cercospora to Avocados  
Product Name: COC WP  
EPA Registration Number: 45002-17  
Application Date: 04-Nov-2014  
Decision Number: 497401

Dear Ms. Miter:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please contact Tony Kish by phone at 703 308-9443, or via email at [kish.tony@epa.gov](mailto:kish.tony@epa.gov); or you may contact Craig Reeves by phone at 703 347-0486, or via email at [reeves.craig@epa.gov](mailto:reeves.craig@epa.gov).

Sincerely,

A handwritten signature in black ink that reads "Tony Kish".

Tony Kish, Product Manager 22  
Fungicide Branch  
Registration Division (7505P)  
Office of Pesticide Programs

2/37

NOTIFICATION

DEC 11 2014

COC WP

Copper Oxychloride Agricultural Fungicide

ACTIVE INGREDIENT:		
Copper Oxychloride*	.....	84.04%
OTHER INGREDIENTS:	.....	15.96%
TOTAL:	.....	100.00%

\*(Metallic copper equivalent 50.0%)  
CAS # 1332-65-6

KEEP OUT OF REACH OF CHILDREN

### CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
IF INHALED	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
IF SWALLOWED	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.</p>	

EPA Reg. No. 45002-17

EPA Est. No. 45002-MX-2

Net Contents:

Manufactured For:  
ALBAUGH, LLC  
ANKENY, IA



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wear long sleeved shirt, long pants, shoes and socks.

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical-resistant to this product are any waterproof material. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

Mixers, loaders, applicators and other handlers must wear the following:

- 1. Long-sleeved shirt and long pants
- 2. Chemical resistant gloves made of any waterproof material
- 3. Shoes plus socks
- 4. Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROL STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- 1. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- 2. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and change into clean clothing.
- 3. Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may contaminate water through runoff. This product has a potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours provide the following instructions are followed.

Not for Greenhouse use.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water, is:

1. Coveralls over long-sleeved shirt and pants
2. Chemical resistant gloves made of any waterproof material
3. Chemical-resistant footwear plus socks
4. Protective eyewear
5. Chemical-resistant headgear for overhead exposure

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests or nurseries.

Do not enter or allow others to enter until sprays have dried.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal of this product.

PESTICIDE STORAGE: Store in a secure, cool, dry area.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Completely empty container into application equipment by shaking and tapping sides and bottom to loosen clinging particles. When completely empty, offer for recycling if available, or dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

### PRODUCT INSTRUCTIONS

COC WP may be applied by aerial, or by dilute or concentrate ground sprayers on crops and at rates given on this label unless specifically prohibited for that crop use. Sufficient spray volume and spray pressure are essential to thoroughly penetrate the plant canopy and give thorough spray coverage at the times indicated. On crops sensitive to copper fungicides use the higher volumes of spray water per acre.

Use the higher dosage rates of COC WP on mature trees, or when disease pressure is severe or weather conditions warrant.

When using adjuvants or other pesticides in combinations with this product, always observe the caution statements on the product's label and required days before harvest. Observe the most restrictive of the labeling limitations and precautions of all products used in mixtures. Sprays of COC WP may be applied up to day of harvest. Residue is exempt from a residue tolerance.

Before mixing with other products in spray tank, be sure that products are compatible. COC WP should not be applied in spray water having a pH of less than 6.5 as phytotoxicity may result. Also avoid using water having a pH of greater than 9.0 as effectiveness may be reduced.

### SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and the method of application (e.g., ground, aerial, airblast, chemigation) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

#### Droplet Size

Apply only as a medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

#### Wind Speed

Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition (approximately 3 to 10 mph), and there are no sensitive areas within 250 feet downwind.

#### Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

#### Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of copper compounds. Where states have more stringent regulations, they must be observed.

#### Equipment

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

#### For aerial application:

The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.

Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety.

When applications are made with a crosswind, the swath must be displaced downwind. The applicator must compensate for this displacement at the up and downwind edge of the application area by adjusting the path of the aircraft upwind.

For groundboom application:

Do not apply with a nozzle height greater than 4 feet above the crop canopy. Handlers may be in the area during application.

MINIMUM RECOMMENDED SPRAY VOLUME

In gallons water per acre (GPA)

If crop is sensitive to copper sprays, higher volumes of spray water will decrease potential injury. A full dilute spray on tree crops means the maximum amount of spray when uniformly applied that an acre of such trees will hold to the point that excess spray begins to drip off. Thus the dilute spray volume per acre will depend on tree size and leaf surface per acre. The following listed dilute spray volumes is the volume that will generally provide such coverage on average size of full leaf trees. A concentrate spray is a spray applied in less volumes than a dilute. The extent of the concentration varies by equipment used. Thus the following spray volumes for a concentrated spray are the minimum volumes recommended per acre.

GROUND SPRAYS - CONCENTRATED AND DILUTE

Citrus -

Concentrate: 100 GPA (Florida 50 GPA).

Dilute: 800 to 1,000 GPA on mature fruit trees and decrease towards 100 GPA as tree size decreases.

Fruit and Nut Trees -

Concentrate: 50 GPA

Dilute spray: 250 GPA for mature fruit trees, and to 400 to 800 GPA for other tree crops depending on size. (On young fruit tree plantings, use a minimum of 15 gallons spray per acre).

Vegetable and Field Crops -

Concentrate: 20 GPA.

Dilute: 100 to 125 GPA

AIR APPLICATION

For vegetables and field crops, use 3 to 20 gallons per acre. For tree crops, use 10 to 20 gallons per acre.

GENERAL CHEMIGATION INSTRUCTIONS

Do not apply this product through any irrigation system using aluminum parts or components as damage to the system may occur. Such application is prohibited regardless of whether the irrigation system is flushed with water after use of this product.

Apply this product only through one or more of the following types of systems: sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveling gun, solid set, or hand move irrigation system(s) which contain no aluminum parts or components. Do not apply this product through any other type of irrigation system.

Crop injury or lack of effectiveness can result from nonuniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide safety device for public water systems is in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

## CHEMIGATION SYSTEMS CONNECTED TO PUBLIC WATER SYSTEMS

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into the reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

When mixing, fill nurse tank half full with water. Add COC WP slowly to tank while hydraulic or mechanical agitation is operating and continue filling with water. Stickers, spreaders, insecticides, nutrients, etc. should be added last. If compatibility is in question, use the compatibility jar test before mixing a whole tank. Because of the wide variety of possible combinations which can be encountered, observe all cautions and limitations on the label of all products used in mixtures.

COC WP should be added through a traveling irrigation system continuously or at the last 30 minutes of solid set or hand moved irrigation systems. Agitation is recommended.

### SPRINKLER CHEMIGATION

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back towards the injection pump.

The pesticide injection pipeline must also contain a functional normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

When mixing, fill nurse tank half full with water. Add COC WP slowly to tank while hydraulic or mechanical agitation is operating and continue filling with water. Stickers, spreaders, insecticides, nutrients, etc. should be added last. If compatibility is in question, use the compatibility jar test before mixing a whole tank. Because of the wide variety of possible combinations which can be encountered, observe all cautions and limitations on the label of all products use in mixtures.

COC WP should be added through a traveling irrigation system continuously or at the last 30 minutes of solid set or hand moved irrigation systems. Agitation is recommended.

MIXING INSTRUCTIONS FOR SPRAY APPLICATION

Fill the spray tank one-fourth to one-third full with clean water. Start agitation (NOTE: Proper agitation creates a rippling or rolling action on the liquid surface). Add COC WP at the recommended rate.

Mix thoroughly and then add enough water to fill spray tank. Maintain sufficient agitation during mixing and during application of sprays to ensure a uniform spray mixture. When tank mixing with other pesticides, add wettable powders or dry flowables first and emulsifiable concentrates or spreader-stickers last. Before adding a second pesticide, be sure that prior product is well mixed and suspended before adding the next ingredient.

RECOMMENDED APPLICATION RATES

FROST INJURY PROTECTION

Bacterial ice nucleation inhibitor - Application of COC WP to all crops listed on this label at rates and stages of growth indicated on this label at least 24 hours and not more than 72 hours prior to anticipated frost conditions, will afford control of ice nucleating bacteria (*Pseudomonas syringae*, *Erwinia herbicola*, and *Pseudomonas fluorescens*) and may thereby provide some protection against light frost. The degree of frost protection will vary with weather conditions and other factors. Not recommended for those geographical areas where weather conditions favor severe frost.



FRUIT, NUT & VINE CROPS

ALMONDS			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Dormant to Pink Bud Season:  Bacterial Blast (Pseudomonas)  Coryneum Blight (Shot hole)	8.0 – 16.0 (4.0 - 8.0 lbs Cu/A)	7	Use at dormant to early pink bud.  A second application in late dormant before foliage buds swell may be necessary when frequent rainfall occurs.  For blast control in sprinkler irrigated orchards or where disease is severe, apply 2-4 sprays or as many as required at 1.0 – 3.0 lbs per acre at 2 week post-bloom intervals or just before sprinkling.  Slight leaf injury may occur from post-bloom spray.
Bloom/Growing Season:  Coryneum Blight Blossom Brown Rot	3.0 (1.5 lbs Cu/A)	5	Use during the early bloom stage (popcorn).  To avoid plant injury, do not use after full bloom.
<b>RESTRICTIONS</b> Maximum single dormant application rate is 16.0 lbs/A (8.0 lbs. metallic copper equivalent) Maximum single bloom/growing application rate is 3.0 lbs/A (1.5 lbs. metallic copper equivalent) Maximum annual application rate is 36.0 lbs/A (18.0 lbs metallic copper equivalent)			



APPLES			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Fall & Late Dormant:  Anthracnose European Canker Pseudomonas Syringae	12.0 – 16.0 (6.0 - 8.0 lbs Cu/A)	Only one dormant application allowed per season	Apply before fall rains.  Use on yellow varieties may cause discoloration. To avoid potential fruit discoloration, pick fruit before spraying foliage.
Fireblight	12.0 (6.0 lbs Cu/A)		Make application between silver-tip and green-tip.  ATTENTION: Phytotoxicity may occur from late application (Discontinue use when green-tip is 1/2 inch.)
Bloom & Growing Season:  Fireblight	1.0 (0.5 lbs Cu/A)	5	Extended spray schedule where fruit finish is not a concern. Continued applications may be made at 5 – 7 day intervals. NOTE: Crop injury may occur from extended spray schedule. Not intended for fresh market apples due to possible russetting. The addition of 1 – 3 lbs of lime per pound of COC WP may reduce injury.
Crown or Collar Rot (Phytophthora cactorum)	N/A	5	Apply either in early spring or in fall after harvest each year.  Do not use if soil pH is below 5.5 or copper toxicity may result.  Mix 0.4 lbs of product in 100 gallon of water. Apply 4 gallons of suspension as a drench on the lower trunk area of each tree.
<b>RESTRICTIONS</b> Maximum single dormant season application rate is 16.0 lbs/A (8.0 lbs. metallic copper equivalent) Maximum single growing season application rate is 1.0 lbs/A (0.5 lbs. metallic copper equivalent) Maximum annual application rate is 32.0 lbs/A (16 lbs. metallic copper equivalent)			

APRICOTS			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Fall & Late Dormant:  Anthracnose European Canker Pseudomonas Syringae	12.0 – 16.0 (6.0 - 8.0 lbs Cu/A)	Only one dormant application allowed per season	Apply before fall rains.  Use the higher rates when conditions favor disease.
Bloom/Growing Season:  Coryneum Blight (Shot Hole) Blossom Brown Rot	2.0 – 3.0 (1.0 – 1.5 lbs Cu/A)	5	Apply at popcorn to full bloom as a full cover spray.  To avoid spray injury, do not apply after full bloom.
<b>RESTRICTIONS</b> Maximum single dormant application rate is 16.0 lbs/A (8.0 lbs. metallic copper equivalent) Maximum single bloom/growing application rate is 3.0 lbs/A (1.5 lbs. metallic copper equivalent) Maximum annual application rate is 36.0 lbs/A (18.0 lbs metallic copper equivalent)			

AVOCADOS			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Anthracnose, Blotch, Scab, <b>Cercospora</b>	4.0 – 6.3 (2.0 - 3.15 lbs Cu/A)	14	Apply when bloom buds begin to swell.  Continue application at 14 – 28 day intervals for 5 to 6 applications.  Use higher rate when conditions favor disease.
<b>RESTRICTIONS</b> Maximum single application rate is 6.3 lbs/A (3.15 lbs metallic copper equivalent) Maximum annual application rate is 37.8 lbs/A (18.9 lbs metallic copper equivalent)			



BANANAS			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Sigatoka (black and yellow)	2.1 (1.05 lbs Cu/A)	7	Apply by air at 2.1 lbs. per acre in 3 gallons of water containing 0.5 gallon agricultural oil. Apply on a 7 - 14 day schedule throughout the wet season.  Apply at 14 - 21 day intervals during dry periods.
Black Pitting	2.1 (1.05 lbs Cu/A)	7	Dilute in 50 - 100 gallons of water and apply directly to the fruit stem and include the basal portion of the leaf crown.  Apply during the first and second weeks after emergence.
<b>RESTRICTIONS</b> Maximum single application rate is 2.1 lbs/A (1.05 lbs metallic copper equivalent) Maximum annual application rate is 37.8 lbs/A (18.9 lbs metallic copper equivalent)			

BRAMBLES (Blackberry, Santiams, Logans, Boysens, Marions, Auroras, Cascades, Chehalems, Rasperry & Thornless Evergreens)			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Anthracnose, Leaf & Cane Spot, Purple Blotch, Yellow Rust	4.0 (2.0 lbs Cu/A)	7	Make fall spray application after harvest.  Apply delayed dormant spray after pruning/training in spring.
	2.0 (1.0 lbs Cu/A)	7	Apply when leaf buds begin to open and repeat when flower buds show white.  NOTE: Crop injury may occur if applied to foliage under hot or moist environmental conditions. Discontinue applications if injury noted.
<b>RESTRICTIONS</b> Maximum single application rate is 4.0/A (2.0 lbs metallic copper equivalent) Maximum annual application rate is 20.0 lbs/A (10.0 lbs metallic copper equivalent)			





CITRUS (Grapefruit, Kumquat, Lemon, Orange, Pummelo, Tangelo, Tangerine & Lime)			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Melanose, Scab, Pink Pitting, Greasy Spot,	4.0 – 6.3 (2.0 – 3.15 lbs Cu/A)	7	Apply as pre-bloom and post-bloom sprays.  Use higher rates when conditions favor disease.
Brown Rot, Septoria Spot	4.0 – 6.3 (2.0 – 3.15 lbs Cu/A)	7	Apply beginning in the fall and continuing as needed.  For Brown Rot, apply to skirts of trees to a height of at least 4 feet.  Apply also to bare ground one foot beyond skirt. Use higher rates when conditions favor disease.  NOTE: In California, in areas subject to copper injury, add 1/4 lb. of high quality lime per lb of COC WP.
Citrus Canker (SUPPRESSION ONLY)	6.3 (3.15 lbs Cu/A)	7	Spraying flushes 7-14 days after shoots begin to grow.  Young fruit may need additional application. Number and timing of applications will depend on disease pressure.  Under heavy disease pressure, each flush of new growth should be sprayed.
Phytophthora Foot Rot	See comment	7	Mix 1.0 lb with one gallon of water and paint trunks of trees from the soil surface to the lowest scaffold limbs.  Apply in May before summer rains and/or in the fall before wrapping trees for freeze protection.  This treatment serves as protection for up to one year, but does not cure existing infections.
Field Nursery Grown To control melanose, scab, pink pitting, greasy spot, brown rot and for citrus canker (suppression).	4.0 – 6.3 (2.0 – 3.15 lbs Cu/A)	7	Apply 2.0 pounds of COC WP per 100 gallons of water.
<b>RESTRICTIONS</b> Maximum single application rate is 6.3 lbs/A (3.15 lbs metallic copper equivalent) Maximum annual application rate is 25.2 lbs/A (12.6 lbs metallic copper equivalent)			

COFFEE			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Coffee Berry Disease (Collectotrichum coffeanum)	3.0 - 4.2 (1.5 - 2.1 lbs Cu/A)	14	Apply after flowering and before the start of long rains and then at 14-28 day intervals until picking.  Use higher rates when rainfall is heavy and disease pressure is high.
Bacterial Blight (Pseudomonas syringae)		14	Begin spray program before the start of long rains and continue until picking.  The critical time of spraying to control disease is just before, during, and after flowering(s), especially when these times coincide with wet weather.  Use higher rates when rainfall is heavy and disease pressure is high.
Iron Spot (Cercospora coffeicola) & Pink Disease (Corticium salmonicolor)	2.0 (1.0 lbs Cu/A)	14	Begin treatment at start of wet season and continue for three applications.
Leaf Rust	3.0 - 4.2 (1.5 - 2.1 lbs Cu/A)	14	Apply before the onset of rain and then at 14 - 21 day intervals while rains continue.  Use higher rates when rainfall is heavy and disease pressure is high.
<b>RESTRICTIONS</b> Maximum single application rate is 4.2 lbs/A (2.1 lbs metallic copper equivalent) Maximum annual application rate is 25.2 lbs/A (12.6 lbs metallic copper equivalent)			





FILBERTS			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Bacterial Blight  (Post Harvest application)	8.0 – 12.0 (4.0 – 6.0 lbs Cu/A)	14	Apply as a post-harvest spray. In seasons of heavy rainfall, apply a second spray when three-quarters of leaves have dropped. Add 1 pint of superior-type oil per 100 gallons of water.
Eastern Filbert Blight			Apply as a dilute spray in adequate water for thorough coverage. Make initial application after harvest in October before heavy winter rains begin. The next application should be made in late February to early March followed by another application 1 month later. If desired, add 1 pint of a sticking agent or superior-type oil per 100 gallons of water.
RESTRICTIONS - Maximum single application rate is 12.0 lbs/A (6.0 lbs metallic copper equivalent) Maximum annual application rate is 48.0 lbs/A (24.0 lbs metallic copper equivalent)			
Note: Permitted only in Washington and Oregon.			

GRAPES			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Black Rot Powdery Mildew Downy Mildew Phomopsis	2.0 – 6.0 (1.0 – 3.0 lbs Cu/A)	3	Begin applications at late dormant up to bud break with subsequent applications throughout the season depending upon disease severity. NOTE: Foliage injury may occur on copper sensitive varieties such as Concord, Delaware, Niagara, and Rosettes. Either test for sensitivity or add 1 to 3 pounds of hydrated lime per pound of COC WP.
RESTRICTIONS Maximum single application rate is 6.0 lbs/A (3.0 lbs metallic copper equivalent) Maximum annual application rate is 40.0 lbs/A (20.0 lbs metallic copper equivalent)			

19/37

GUAVA			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Anthracnose Red Algae	2.5 (1.25 lbs Cu/A)	7	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage.
<b>RESTRICTIONS</b> Maximum single application rate is 2.5 lbs/A (1.23 lbs metallic copper equivalent) Maximum annual application rate is 9.8 lbs product/A (4.92 lbs metallic copper equivalent)			

HOPS			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Downy Mildew	1.0 (.50 lbs Cu/A)	10	Apply as a fungicide crown treatment (after pruning, but before training) as needed.  After training, additional fungicide treatments are needed at 10 day intervals.  Discontinue use 2 weeks before harvest.
<b>RESTRICTIONS</b> Maximum single application rate is 1.0 lb/A (0.5 lbs metallic copper equivalent) Maximum annual application rate is 5.0 lbs/A (2.5 lbs metallic copper equivalent)			

KIWI			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Pseudomonas syringae Erwinia herbicola Pseudomonas fluorescens	4.2 (2.1 lbs Cu/A)	30	Make applications on a monthly basis. A maximum of 3 applications may be made.
<b>RESTRICTIONS</b> Maximum single application rate is 4.2 lbs/A (2.1 lbs metallic copper equivalent) Maximum annual application rate is 12.6 lbs/A (6.3 lbs metallic copper equivalent)			

LITCHI			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Anthracnose	2.4 (1.2 lbs Cu/A)	7	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage.
<b>RESTRICTIONS</b> Maximum single application rate is 2.4 lbs/A (1.2 lbs metallic copper equivalent) Maximum annual application rate is 9.6 lbs/A (4.8 lbs metallic copper equivalent)			

MACADAMIA NUTS			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Anthracnose	4.7 (2.35 lbs Cu/A)	7	Initiate sprays at first sign of flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage
Blossom blight Raceme blight	3.0 – 4.0 (1.5 – 2.0 lbs Cu/A)	7	Apply during peak raceme development and bloom period. Use higher rates when conditions favor disease.
<b>RESTRICTIONS</b> Maximum single application rate is 4.7 lbs/A (2.36 lbs metallic copper equivalent) Maximum annual application rate is 18.8 lbs/A (9.44 lbs metallic copper equivalent)			

MAMEY SAPOTE			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Anthracnose Algal Leaf Spot	3.0 – 4.2 (1.5 - 2.1 lbs Cu/A)	14	Apply when conditions favor disease development.  Repeat on 14-30 day schedule as disease severity and environmental conditions dictate. Use higher rates when conditions favor disease.
<b>RESTRICTIONS</b> Maximum single application rate is 4.2 lbs/A (2.1 lbs metallic copper equivalent) Maximum annual application rate is 16.8 lbs/A (8.4 lbs metallic copper equivalent)			

MANGO			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Anthracnose	4.0 – 5.2 (2.0 – 2.6 lbs Cu/A)	30	Apply monthly after fruit set until harvest.
<b>RESTRICTIONS</b> Maximum single application rate is 5.2 lbs/A (2.6 lbs metallic copper equivalent) Maximum annual application rate is 36.4 lbs/A (18.2 lbs metallic copper equivalent)			

OLIVES			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Peacock Spot Olive Knot	5.0 – 6.3 (2.5 – 3.15 lbs Cu/A)	30	Apply before winter rains fall.  A second application in early spring should be made if disease is severe.  Use higher rates when conditions favor disease.
<b>RESTRICTIONS</b> Maximum single application rate is 6.3 lbs/A (3.15 lbs metallic copper equivalent) Maximum annual application rate is 12.6 lbs/A (6.3 lbs metallic copper equivalent)			

PAPAYA			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Anthracnose	4.0 – 5.2 (2.0 – 2.6 lbs Cu/A)	14	Begin application before disease is expected to appear.  Repeat at 14 day intervals. Use the higher rates when conditions favor disease.  The addition of a suitable spreader-sticker may be desirable especially during periods of heavy rains.
<b>RESTRICTIONS</b> Maximum single application rate is 5.2 lbs/A (2.63 lbs metallic copper equivalent) Maximum annual application rate is 42.4 lbs/A (21.2 lbs metallic copper equivalent)			



PEARS, QUINCE			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Bloom & Growing Season:  Fireblight	1.0 (0.5 lbs Cu/A)	5	Apply at 5 day intervals throughout bloom period. Excessive dosages may cause fruit russet.
Fall & Late Dormant Season:  Pseudomonas blight	12.0 – 16.0 (6.0 – 8.0 lbs Cu/A)	Only one dormant application allowed per season	Apply before fall rain begins.
<b>RESTRICTIONS</b> Maximum single dormant season application rate is 16.0 lbs/A (8.0 lbs metallic copper equivalent) Maximum single growing season application rate is 1.0 lb/A (0.5 lbs metallic copper equivalent) Maximum annual application rate is 32.0 lbs/A (16.0 lbs metallic copper equivalent)			

PECANS			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Shuck and Kernel rot (Phytophthora cactorum)  Zonate leaf spot (Cristulariella pyramidalis)	2.0 – 4.2 (1.0 – 2.1 lbs Cu/A)	14	Apply in sufficient water for good coverage at 2-4 week intervals starting at kernel growth and continuing until shucks open.  Use the higher rate and shorter intervals if frequent rainfall occurs.
Mosses Algae Lichen	See Comment	Make only one application per year	Mix 2 lbs per 100 gallons spray plus spreader-sticker on a dilute spray basis and apply in dormant season before buds swell, thoroughly wetting limbs and mosses.
<b>RESTRICTIONS</b> Maximum single application rate is 4.2 lbs/A (2.1 lbs metallic copper equivalent) Maximum annual application rate is 16.4/A (8.4 lbs metallic copper equivalent)			



PISTACHIOS			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Botrytis Blight, Botryosphaeria Panicle, Shoot Blight, Septoria Leaf Blight, Late Blight (Alternaria)	3.0 – 4.2 (1.5 - 2.1 lbs Cu/A)	14	Make initial application at bud swell and repeat on a 14 - 28 day schedule.  Use higher rates when conditions favor disease.
<b>RESTRICTIONS</b> Maximum single application rate is 4.2 lbs/A (2.1 lbs metallic copper equivalent) Maximum annual application rate is 16.8 lbs/A (8.4 lbs metallic copper equivalent)			

PLUMS & PRUNES			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Dormant Season:  Coryneum blight (Shot hole)	8.0 – 16.0 (4.0 – 8.0 lbs Cu/A)	7	Apply as a dormant spray. Use the higher rate when rainfall is heavy and/or disease pressure is high.
Bloom & Growing Season:  Brown rot blossom blight, Black Knot	3.0 (1.5 lbs Cu/A)	5	Apply full cover application at pink, red or early white bud stage. Use the higher rate when disease pressure is heavy or conditions favor disease development.
<b>RESTRICTIONS</b> Maximum single dormant, late dormant and pink bud application rate is 16.0 lbs/A (8.0 lbs metallic copper equivalent) Maximum single growing season application rate is 3.0 lbs/A (1.5 lbs metallic copper equivalent) Maximum annual application rate is 36.0 lbs/A (18.0 lbs metallic copper equivalent)			

STRAWBERRIES			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Downy Mildew Leaf Spot Leaf Blight	2.0 - 3.0 (1.0 – 1.5 lbs Cu/A)	7	Begin application when plants are established and continue on a weekly schedule throughout season. Apply in at least 20 gallons of water. Use the higher rates when conditions favor disease.  NOTE: Discontinue applications if signs of phytotoxicity appear.
<b>RESTRICTIONS</b> Maximum single application rate is 3.0 lbs/A (1.5 lbs metallic copper equivalent) Maximum annual application rate is 16.0 lbs/A (8.0 lbs metallic copper equivalent)			



26/37

VEGETABLE AND FIELD CROPS

ALFALFA			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM RETREATMENT INTERVAL	COMMENT
Cercospora & Leptosphaerulina Leaf Spots	1.0 (0.5 lbs Cu/A)	30 Days	Apply 10 to 14 days before each harvest or earlier if disease threatens.  Apply with ground or aerial equipment.  Spray injury may occur with sensitive varieties such as Lahontan.
<b>RESTRICTIONS:</b> Maximum single application rate is 1.0 lb/A (0.5 lbs metallic copper equivalent) Maximum annual application rate is 2.0 lb/A (1.0 lbs metallic copper equivalent)			

BEANS (Dry, Green)			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Bacterial Blight (Halo & Common)  Brown spot	1.0 - 1.5 (0.5 - .75 lbs Cu/A)	7	For protective sprays apply first application when plants are five to six inches high. Apply on 7 - 14 day schedule depending on local conditions. Use higher rate for more severe disease pressure.
<b>RESTRICTIONS</b> Maximum single application rate is 1.5 lbs/A (0.75 lbs. metallic copper equivalent) Maximum annual application rate is 9.0 lbs/A (4.5 lbs metallic copper equivalent)			

SUGAR BEETS & TABLE BEETS			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Cercospora Leaf Spot	2.0 - 2.5 (1.0 - 1.25 lbs Cu/A)	10	Begin applications when conditions first favor disease development and repeat at 10 to 14 day intervals as needed. Use the higher rate when disease is severe.
<b>RESTRICTIONS</b> Maximum single application rate is 2.5 lbs/A (1.25 lbs metallic copper equivalent) Maximum annual application rate is 15.0 lbs/A (7.5 lbs metallic copper equivalent)			

CARROTS			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Carrot Blight (Cercospora)	2.0 (1.0 lbs Cu/A)	7	Begin application when disease first threatens and repeat at 7 to 14 day intervals as needed depending on disease severity.
<b>RESTRICTIONS</b> Maximum single application rate is 2.0 lbs/A (1.0 lbs metallic copper equivalent) Maximum annual application rate is 10.0/A (5.0 lbs metallic copper equivalent)			

CUCURBITS (CUCUMBERS, CANTALOUPE, HONEYDEWS, MUSKMELONS, PUMPKINS, SQUASH & WATERMELONS)			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Alternaria Leaf Spot Angular Leaf Spot Anthracnose Downy Mildew Powdery Mildew Gummy Stem Blight Watermelon Bacterial Fruit Blotch (suppression)	2.0 (1.0 lbs Cu/A)	5	Begin application when conditions are favorable for disease development. Repeat at 5-10 day intervals.  Use higher rates when conditions favor disease.  NOTE: Crop injury may occur from application at higher rates and shorter intervals.  Discontinue use if injury occurs.
<b>RESTRICTIONS</b> Maximum single application rate is 2.0 lbs/A (1.0 lbs metallic copper equivalent) Maximum annual application rate is 10.5 lbs/A (5.25 lbs metallic copper equivalent)			

CELERY & CELERIAC			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Early, Late & Bacterial Blights	2.0 (1.0 lbs Cu/A)	7	Apply as soon as plants are first established in the field, then every 7 days depending on severity and weather.
<b>RESTRICTIONS</b> Maximum single application rate is 2.0 lbs/A (1.0 lbs metallic copper equivalent) Maximum annual application rate is 10.0 lbs/A (5.0 lbs metallic copper equivalent)			

CHIVES			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Downy Mildew	1.0 (0.5 lbs Cu/A)	7	Begin applications when plants are established in the field. Repeat applications every 7-10 days as dictated by disease conditions.
<b>RESTRICTIONS</b> Maximum single application rate is 1.0 lb/A (0.5 lbs metallic copper equivalent) Maximum annual application rate is 5.0 lbs/A (2.5 lbs metallic copper equivalent)			

CRUCIFERS (Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collard Greens, Mustard Greens, & Turnip Greens)			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Downy Mildew Black Rot (Xanthomonas) Black Leaf Spot (Alternaria)	1.0 (0.5 lbs Cu/A)	7	Begin application after transplants are set in the field, or shortly after emergence of field seeded crops or when conditions favor disease development.  (CAUTION: A slight reddening of older leaves may occur on broccoli, and a slight flecking of wrapper leaves may occur on cabbage.)
<b>RESTRICTIONS</b> Maximum single application rate is 1.0 lbs/A (0.5 lbs metallic copper equivalent) Maximum annual application rate is 5.0 lbs/A (2.5 lbs metallic copper equivalent)			

DILL			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Phoma Leaf Spot, Rhizoctonia Foliage Blight	1.5 (0.75 lbs Cu/A)	7	Begin applications when plants are first established in the field and repeat at 7-10 day intervals depending upon disease severity and environmental conditions.
<b>RESTRICTIONS</b> Maximum single application rate is 1.5 lbs/A (0.75 lbs metallic copper equivalent) Maximum annual application rate is 7.5 lbs/A (3.75 lbs. metallic copper equivalent)			



30/37

**LETTUCE, ENDIVE & ESCAROLE**

DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Downy Mildew	1.0 – 2.0 (0.5 - 1.0 lbs Cu/A)	5	Begin treatment when disease first appears and repeat every 5 - 10 days as needed to suppress disease.

NOTE: Flecking and/or yellowing of leaves will occur under certain environmental conditions such as extended periods of moist weather, acid rains, or other conditions favoring reduced pH on leaf surfaces. Injury may be severe enough to reduce crop value. Increasing the volume of spray water may decrease phytotoxicity potential.

**RESTRICTIONS**

Maximum single application rate is 2.0 lbs/A (1.0 lb metallic copper equivalent)  
 Maximum annual application rate is 16.0 lbs/A (8.0 lbs metallic copper equivalent)

**ONION & GARLIC**

DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Purple Blotch Downy Mildew	2.0 (1.0 lbs Cu/A)	7	Begin when plants are 4 to 6 inches high and repeat at 7 to 10 day intervals as needed depending upon disease pressure. Can cause phytotoxicity to leaves.
Bacterial Blight	1.0 – 1.5 (0.5 - .75 lbs Cu/A)		

**RESTRICTIONS**

Maximum single application rate is 2.0 lbs/A (1.0 lb metallic copper equivalent)  
 Maximum annual application rate is 12.0 lbs/A (6.0 lbs metallic copper equivalent)

**PARSLEY**

DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Bacterial Blight (Pseudomonas sp.)	2.0 (1.0 lbs Cu/A)	10	Begin applications when plants are first established in the field and repeat at 10 day or greater intervals depending upon disease severity and environmental conditions.

**RESTRICTIONS**

Maximum single application rate is 2.0 lbs/A (1.0 lbs metallic copper equivalent)  
 Maximum annual application rate is 4.0 lbs/A (2.0 lbs metallic copper equivalent)

PEANUTS			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Cercospora Leaf Spot	1.5 (0.75 lbs Cu/A)	7	Begin spraying 25-40 days after planting or when disease symptoms appear.  Use sufficient water to get adequate coverage.  Continue applications at 7 to 14 day intervals. Reduce spray interval to 7 days during humid weather.
<b>RESTRICTIONS</b> Maximum single application rate is 1.5 lbs/A (0.75 lbs metallic copper equivalent) Maximum annual application rate is 9.0 lbs/A (4.5 metallic copper equivalent)			

PEAS			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Powdery Mildew	1.5 (0.75 lbs Cu/A)	7	Begin spray treatment when disease symptoms first appear.  Repeat applications at weekly intervals.
<b>RESTRICTIONS</b> Maximum single application rate is 1.5 lbs/A (0.75 lbs metallic copper equivalent) Maximum annual application rate is 7.5 lbs/A (3.75 lbs metallic copper equivalent)			

PEPPERS			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Bacterial Spot	1.5 (0.75 lbs Cu/A)	3	Apply, when disease threatens, in sufficient water to provide adequate coverage.  Use at 3 to 14 day intervals depending on disease severity.
<b>RESTRICTIONS</b> Maximum single application rate is 1.5 lbs/A (0.75 lbs metallic copper equivalent) Maximum annual application rate is 23.5 lbs/A (11.75 lbs metallic copper equivalent)			



TOMATOES (Processing)			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Early Blight Bacterial Speck Bacterial Spot Anthracnose Gray Leaf Mold Gray Leaf Spot Septoria Leaf Spot Late Blight	1.0 (0.5 lbs Cu/A)	3	When disease threatens, apply at 7 - 10 day intervals, more frequently when disease is severe.
<b>RESTRICTIONS</b> Maximum single application rate is 1.0 lb/A (0.5 lbs metallic copper equivalent) Maximum annual application rate is 34.5 lbs/A (17.25 lbs metallic copper equivalent)			

TOMATOES (Fresh Market)			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Early Blight Bacterial Speck Bacterial Spot Anthracnose Gray Leaf Mold Gray Leaf Spot Septoria Leaf Spot Late Blight	2.0 - 3.2 (1.0 - 1.6 lbs Cu/A)	3	When disease threatens, apply at 7 - 10 day intervals, more frequently when disease is severe.
<b>RESTRICTIONS</b> Maximum single application rate is 3.2 lb/A (1.6 lbs metallic copper equivalent) Maximum annual application rate is 16.0 lbs/A (8.0 lbs metallic copper equivalent)			

WATERCRESS			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Cercospora Leaf Spot	1.0 (0.5 lbs Cu/A)	7	Begin application when plants are first established in the field, repeating at 7-14 day intervals depending on disease severity and environmental conditions. Do not exceed 4 applications per crop. Apply using ground spray equipment at no less than 50 gallons of spray solution per acre.
<b>RESTRICTIONS</b> Maximum single application rate is 1.0 lb/A (0.5 lbs metallic copper equivalent) Maximum annual application rate is 4.0 lbs/A (2.0 lbs metallic copper equivalent)			

WHEAT, BARLEY & OATS			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Septoria Leaf Blotch Helminthosporium Spot Blotch	1.0 (0.5 lbs Cu/A)	10	Make first application at early heading and follow with second application 10 days later.
<b>RESTRICTIONS</b> Maximum single application rate is 1.0 lb/A (0.50 lbs metallic copper equivalent) Maximum annual application rate is 2.0 lbs/A (1.0 lbs metallic copper equivalent)			

ORNAMENTAL PLANTS, HEDGES, AND FOREST TREES

DOUGLAS FIR			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Rhabdocline needlecast	2.0 – 4.0 (1.0 – 2.0 lbs Cu/A)	7	Begin applications at bud break and repeat at 7 – 28 day intervals. Use higher rates when conditions favor disease.
<b>RESTRICTIONS</b> Maximum single application rate is 4.0 lbs/A (2.0 lbs metallic copper equivalent) Maximum annual application rate is 40.0 lbs/A (20.0 lbs metallic copper equivalent)			

LIVE OAK			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Ball Moss	Mix 3.0 – 4.0 lbs per 100 gallons of water.  (1.5 - 2.0 lbs Cu)	A second application may be required after 12 months.	Apply in spring after heavy rain, using 1.5 gallons of spray per foot of tree height. Make sure to wet tufts thoroughly.  NOTE: COC WP may be injurious to some ornamentals grown under live oaks.  This product may be reactive on metal and masonry surfaces such as galvanized roofing. Avoid contact with metal surfaces. Do not spray on cars, houses, lawn furniture, etc.
<b>RESTRICTIONS</b> Maximum single application rate is 4.0 lbs/A (2.0 lbs metallic copper equivalent) Maximum annual application rate is 40.0 lbs/A (20.0 lbs metallic copper equivalent)			

SYCAMORE			
DISEASE	APPLICATION RATE (lbs product/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Anthracnose	2.0 – 4.0 (1.0 – 2.0 lbs Cu/A)	7	Make two applications as a full cover spray.  Use a minimum of 100 gallons water per acre.  Make first application at bud crack and second application 7 to 14 days later at 10% leaf expansion.
<b>RESTRICTIONS</b> Maximum single application rate is 4.0 lbs/A (2.0 lbs metallic copper equivalent) Maximum annual application rate is 40.0 lbs/A (20.0 lbs metallic copper equivalent)			

TURFGRASS			
DISEASE	APPLICATION RATE (pints)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Algae	4.0 – 6.0 (2.0 – 3.0 lbs Cu/A)	10	May be used as a maintenance spray as needed. May be used alone or in combination with fungicides such as dithiocarbamates.  Use a minimum of 100 gallons of water per acre.  Phytotoxicity may depend on varietal differences. Apply the recommended rate to a small area and observe 7 - 10 days for phytotoxicity. If phytotoxicity occurs, discontinue use.
<b>RESTRICTIONS</b> Maximum single application rate is 6.0 lbs/A (3.0 lbs metallic copper equivalent) Maximum annual application rate is 18.0 lbs/A (9.0 lbs metallic copper equivalent)			

ORNAMENTALS

Notice to User: Plant sensitivities to COC WP have been found to be acceptable in specific genera and species listed on this label; however, phytotoxicity may occur. Due to the large number of species and varieties of ornamentals and nursery plants, it is impossible to test every one for sensitivity to COC WP. Neither the manufacturer nor seller has determined whether or not COC WP can be safely used on ornamental or nursery plants not listed on this label. The user should determine if COC WP can be used safely prior to commercial use. In a small area, apply the recommended rates to the plants in question, i.e., bedding plants, foliage, etc., and observe for 7-10 days for symptoms of phytotoxicity prior to commercial use.

Use COC WP on container, bench, or bed-grown ornamentals in greenhouses or outdoor nurseries, for professional use on ornamentals grown for indoor and outdoor landscaping, and for control of bacterial and fungal diseases of foliage, flowers and stems.

Apply as a thorough coverage spray using 1.0 lb per 100 gallons of water. Begin application at first sign of disease and repeat at 7 to 14 day intervals as needed; use shorter interval during periods of frequent rains or when severe disease conditions persist.

COC WP may be used as a maintenance spray alone or in combination with other fungicides such as the dithiocarbamates.

Restrictions:

- Maximum single application rate is 4.0 lbs/A (2.0 lbs metallic copper equivalent)
- Maximum annual application rate is 40.0 lbs/A (20 lbs metallic copper equivalent)

ORNAMENTAL/DISEASES:

- Althea (Rose of Sharon)/Bacterial Leaf Spot
- Aralia/Xanthomonas & Cercospora Leaf Spots, Alternaria
- Arborvitae/Alternaria Twig Blight, Cercospora Leaf Blight
- Azalea\*/Cercospora Leaf Spot, Botrytis Blight, Phytophthora Dieback & Powdery Mildew
- Begonia/Xanthomonas Leaf Spot, Anthracnose
- Bougainvillea/Anthracnose, Bacterial Leaf Spot
- Bulbs (Easter Lily\*\*, Tulip, Gladiolus)/Anthracnose, Botrytis Blight
- Camellia/Anthracnose, Bacterial Leaf Spot
- Camphor Tree/Pseudomonas Leaf Spot
- Canna/Pseudomonas Leaf Spot
- Carnation\*/Alternaria Blight, Pseudomonas Leaf Spot, & Botrytis Blight
- Chinese Tallow Tree/Bacterial Leaf Spot (Xanthomonas sp., Pseudomonas sp.)
- Chrysanthemum\*/Septoria Leaf Spot & Botrytis Blight
- Cotoneaster/Botrytis Blight
- Dahlia/Alternaria Leaf Spot, Botrytis Gray Mold, Cercospora Leaf Spot
- Date Palm/Pestalotia Leaf Spot
- Dianthus/Bacterial Spot, Bacterial Soft Rot
- Dogwood/Anthracnose
- Dusty Miller/Bacterial Leaf Spot (Pseudomonas cichorii)
- Echinacea/Bacterial Leaf Spot (Pseudomonas cichorii)
- Elm "Drake"/Xanthomonas Leaf Spot
- Euonymus/Botrytis Blight & Anthracnose
- European Fan Palm/Pestalotia Leaf Spot
- Gardenia/Alternaria Leaf Spot, Botrytis Bud Rot, Cercospora Leaf Spot
- Geranium/Alternaria Leaf Spot, Botrytis Gray Mold, Cercospora Leaf Spot
- Gladiolus/Alternaria Leaf Spot, Botrytis Gray Mold, Bacterial Leaf Blight
- Golden Rain Tree/Bacterial Leaf Spot
- Hibiscus/Bacterial Leaf Spot
- Holly Fern/Pseudomonas Leaf Spot
- Impatiens/Bacterial Leaf Spot
- India hawthorn (greenhouse)/Entomosporium Leaf Spot
- Ivy\*/Xanthomonas Leaf Spot
- Ixora/Xanthomonas Leaf Spot
- Juniper (Eastern Red Cedar)/Anthracnose
- Lantana/Bacterial Leaf Spot
- Lilac/Cercospora Leaf Spot
- Loblolly Bay/Anthracnose



- Loquat/Entomosporium maculata, Colletotrichum sp.
- Magnolia (Southern)/Algal Leaf Spot, Anthracnose, Bacterial Leaf Spot
- Mandevillas/Anthracnose
- Marigold/Alternaria Leaf Spot, Botrytis Leaf and Flower Rot, Cercospora Leaf Spot
- Mulberry, Weeping/Bacterial Leaf Spot
- Oak, Laurel/Algal Leaf Spot (Cephaleuros virescens)
- Oleander/Bacterial Leaf Spot, Fungal Leaf Spot
- Pachysandra/Volutella Leaf Blight
- Pansy/Downy Mildew
- Pear (Flowering)/Fireblight, Leaf Spot
- Pentas (Egyptian Star)/Bacterial Leaf Spot (Xanthomonas sp.)
- Peony/Botrytis Blight
- Periwinkle/Phomopsis Stem Blight
- Philodendron/Bacterial Leaf Spot
- Phlox/Alternaria Leaf Spot
- Photinia (Red Tip)/Anthracnose, Entomosporium Leaf Spot
- Pistachio/Anthracnose
- Plantain Lily/Bacterial Leaf Spot
- Powder Puff Plant/Bacterial Leaf Spot
- Pyracantha/Fireblight & Scab
- Queen Palm/Exosporium Leaf Spot, Phytophthora Bud Rot
- Rhododendron/Alternaria Flower Spot
- Rose\*/Powdery Mildew, Black Spot
- Verbena/Xanthomonas Leaf Spot
- Viburnum/Anthracnose
- Washingtonia Palm/Pestalotia Leaf Spot
- Weeping Willow/Anthracnose
- Yucca (Adams Needle)/Cercospora & Septoria Leaf Spot

\*Discoloration of foliage and/or blooms have been noted on some varieties. To prevent residues on commercial plants, do not spray just before selling season.

\*\*For Easter Lily, use 2.0 – 3.0 pints per 100 gallons.

Restrictions (Easter Lily):

- Maximum single application rate is 5.0 lbs/A (2.5 lbs metallic copper equivalent)
- Maximum annual application rate is 150 lbs/A (75 lbs metallic copper equivalent)

NOTICE: To the extent consistent with applicable law, seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use; but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions not reasonably foreseeable to seller; the buyer assumes the risk of any such use.

